

		令和02年度						令和03年度						令和04年度						令和05年度			令和06年度		
		5/20	7/22	11/12	1/18	平均	5/26	7/30	11/26	1/17	平均	5/18	7/20	11/14	1/19	平均	8/21	1/16	平均	9/26	12/13	平均			
中海流入河川調査	大正川 (平成橋)	気温	15.8	25.8	16.0	2.7		24.8	32.5	18.0	7.0		25.7	32.0	16.3	7.3		35.2	6.0						
		水温	18.4	23.6	16.7	8.6		20.7	25.6	13.7	8.6		20.0	27.7	17.7	9.7		28.0	10.2						
		化学的酸素要求量(～R5)	mg/L	3.2	2.8	3.2	5.0	3.6	3.7	3.7	3.6	3.7	4.1	6.9	3.8	4.1	4.7	3.4	4.3	3.9					
		生物化学的酸素要求量(R6)	mg/L	0.66	0.64	1.0	2.4	1.2	0.85	1.00	1.0	1.2	1.0	1.70	1.70	1.0	1.7	1.5	0.83	1.60	1.2				
		りん含有量	mg/L	0.033	0.043	0.090	0.093	0.06	0.052	0.092	0.060	0.085	0.07	0.092	0.230	0.091	0.073	0.12	0.070	0.058	0.06				
	米川 (弥生橋)	水温	0.48	0.74	0.07	0.32	0.40	0.39	0.04	0.09	0.05	0.14	0.09	0.18	0.11	0.06	0.11	0.19	0.20	0.20					
		気温	16.0	24.5	15.0	1.8		24.5	33.0	18.2	8.3		23.5	29.2	18.2	7.2		34.7	6.1		29.0	10.0			
		水温	19.0	23.5	15.3	7.7		20.1	26.6	14.4	8.6		20.0	27.7	15.3	6.4		27.3	9.1		24.6	10.5			
		化学的酸素要求量(～R5)	mg/L	5.1	3.0	5.5	5.1	4.7	5.2	3.5	4.1	4.6	4.4	5.0	13.0	3.6	6.3	3.4	4.9	4.2	0.9	1.5	1.2		
		生物化学的酸素要求量(R6)	mg/L	0.75	0.93	2.4	2.6	1.7	0.83	0.84	1.5	1.6	1.2	0.95	2.20	1.1	1.8	1.5	0.70	4.90	2.8	0.60	1.40	1.0	
境港市内河川調査	油田川 (外江町2061付近)	葉素含有量	mg/L	0.072	0.073	0.34	0.11	0.15	0.100	0.073	0.052	0.088	0.08	0.100	0.290	0.093	0.110	0.15	0.100	1.300	0.70	0.082	0.079	0.08	
		りん含有量	mg/L	0.53	0.69	0.06	0.22	0.38	0.37	0.60	0.20	0.33	0.38	0.14	0.42	0.36	0.28	0.30	0.27	0.14	0.20	0.17			
		水温	18.5	26.3	12.5	1.0		22.6	32.0	16.2	5.8		18.3	29.0	17.2	9.0		35.0	10.0		28.0	10.3			
		気温	17.1	20.2	16.2	11.8		17.6	26.9	15.9	10.2		18.0	22.2	16.5	11.1		23.6	7.5		22.5	14.2			
		化学的酸素要求量(～R5)	mg/L	11.0	5.9	11.0	7.0	8.7	15.0	3.0	8.4	7.3	8.4	22.0	8.8	9.1	8.9	12.2	9.6	6.0	7.8	3.4	3.5	3.5	
	八幡川 (渡町3060-5付近)	生物化学的酸素要求量(R6)	mg/L	6.90	3.2	5.8	3.5	4.9	11.00	2.6	6.8	6.6	6.8	11.00	4.9	4.8	5.0	6.4	5.10	3.9	4.5	3.60	3.3	3.5	
		葉素含有量	mg/L	0.810	0.30	0.97	0.29	0.59	2.800	0.23	0.71	0.73	1.12	1.800	0.57	0.57	0.49	0.86	0.880	0.38	0.63	0.580	0.28	0.43	
		りん含有量	mg/L	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01		
		水温	17.5	26.8	12.8	1.2		24.6	32.1	15.8	5.3		20.7	29.3	16.7	7.6		34.3	5.3		28.0	9.0			
		気温	18.3	22.3	13.1	10.2		21.9	25.6	15.0	10.1		20.1	26.2	15.4	11.8		27.9	11.7		24.3	14.2			
新屋上ノ川	新屋上ノ川	化学的酸素要求量(～R5)	mg/L	13.0	6.1	11.0	5.6	8.9	13.0	5.2	7.4	4.3	7.5	19.0	8.7	5.8	4.5	9.5	5.7	4.4	5.1	3.0	1.8	2.4	
		生物化学的酸素要求量(R6)	mg/L	11.00	4.6	9.0	5.2	7.5	16.00	3.7	5.0	3.3	7.0	9.10	6.4	3.1	3.6	5.6	2.90	3.8	3.4	3.00	3.2	3.1	
		葉素含有量	mg/L	1.000	0.35	0.89	0.25	0.62	1.100	0.24	0.35	0.17	0.47	1.800	0.83	0.31	0.11	0.76	0.360	0.12	0.24	0.460	0.12	0.29	
		りん含有量	mg/L	< 0.01	0.02	< 0.01	0.03	0.03	< 0.01	0.02	< 0.01	< 0.01	0.02	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01		
		水温	14.8	27.7	7.8	0.2		21.8	31.0	11.5	6.1		18.8	27.5	13.0	5.0		32.6	3.8		29.0	7.0			
	才仏川	気温	18.6	24.4	13.9	9.8		17.9	23.7	13.4	5.7		17.7	22.9	15.3	9.1		23.8	8.7		25.0	11.5			
		化学的酸素要求量(～R5)	mg/L	2.7	3.4	3.5	4.3	3.5	2.7	3.1	4.3	3.3	3.4	3.1	5.5	3.5	4.7	4.2	3.4	3.9	3.7	0.8	0.7	0.8	
		生物化学的酸素要求量(R6)	mg/L	0.53	1.4	1.4	3.9	1.8	0.63	1.2	2.4	1.8	1.5	1.10	2.2	1.2	3.1	1.9	0.86	2.5	1.7	0.52	2.6	1.6	
		りん含有量	mg/L	0.047	0.37	0.36	0.58	0.34	0.110	0.21	0.53	0.24	0.27	0.220	0.49	0.22	0.42	0.34	0.089	0.39	0.24	0.120	0.51	0.32	
		水温	15.4	26.8	8.6	(0.6)		21.0	31.6	11.3	6.0		21.4	27.5	13.5	5.2		30.8	3.6		29.5	7.8			
下ノ川	中野承水路	化学的酸素要求量(～R5)	mg/L	3.4	5.7	3.9	4.7	4.4	5.0	5.7	3.8	3.6	4.5	4.1	3.6	3.6	3.5	3.7	3.9	3.7	3.8	0.8	0.6	0.7	
		生物化学的酸素要求量(R6)	mg/L	0.88	0.92	1.3	1.2	1.1	1.10	1.00	0.88	0.95	1.0	1.10	1.00	0.99	0.93	1.0	1.20	0.94	1.1	0.89	0.84	0.9	
		葉素含有量	mg/L	0.120	0.27	0.30	0.12	0.20	0.220	0.27	0.19	0.11	0.20	0.290	0.33	0.21	0.13	0.24	0.380	0.15	0.27	0.350	0.18	0.27	
		りん含有量	mg/L	18.4	28.0	11.0	(0.2)		22.1	31.6	12.8	6.3		21.5	28.2	13.7	6.1		32.0	4.0					
		水温	17.8	22.7	16.8	8.9		19.0	27.1	15.1	10.2		19.5	25.9	16.8	10.9		32.8	10.8						
	下ノ川	化学的酸素要求量(～R5)	mg/L	2.4	2.9	2.2	3.4	2.7	3.8	3.3	2.8	3.4	2.5	4.0	4.8	2.7	3.5	2.9	3.2	3.1					
		生物化学的酸素要求量(R6)	mg/L	0.72	0.85	0.99	1.3	1.0	1.50	0.92	1.30	1.4	1.3	0.86	1.10	1.20	1.0	0.89	0.74	0.8					
		葉素含有量	mg/L	0.150	0.32	0.30	0.22	0.25	0.800	0.32	0.42	0.27	0.45	0.180	0.34	0.13	0.17	0.21	0.420	0.14	0.28				
		りん含有量	mg/L	15.4	27.2	12.0	9.6		23.0	27.8	14.6	9.4		23.0	24.0	15.0	10.7		31.4	10.5		25.5	14.0		
		水温	17.6	22.5	16.1	7.6		18.4	29.8	14.6	10.1		19.7	26.7	17.9	7.9		29.4	10.8						
卸売市場岸壁	下ノ川	化学的酸素要求量(～R5)	mg/L	5.2	5.1	4.5	5.6	5.1	6.2	7.3	5.8	6.3	6.4	8.6	12.0	5.5	6.1	8.1	5.5	5.7	5.6	1.2	1.3	1.3	
		生物化学的酸素要求量(R6)	mg/L	0.97	1.20	1.30	1.80	1.32	0.97	1.20	1.50	1.60	1.32	1.30	1.70	0.87	1.60	1.37	0.44	1.30	0.87	0.27	1.20	0.74	
		葉素含有量	mg/L	0.130	0.15	0.10	0.13	0.13	0.170	0.21	0.13	0.14	0.16	0.410	0.34	0.12	0.13	0.25	0.150	0.13	0.14	0.100	0.20	0.15	
		りん含有量	mg/L	17.9	28.1	12.0	0.8		23.0	32.0	12.8	6.5		21.4	27.0	15.3	7.0		30.8	4.8					
		水温	19.0	25.8	16.2	7.6		18.4	29.8	14.6	10.1		19.7	26.7	17.9	7.9		29.4	10.8						
	渡港	化学的酸素要求量(～R5)	mg/L	2.0	2.5	2.1	2.5	2.3	1.6	3.9	2.8	2.6	2.7	2.0	3.5	2.2	3.0	2.7	2.4	1.5	2.0				
		生物化学的酸素要求量(R6)	mg/L	0.21	0.28	0.28	0.39	0.3	0.27	0.39	0.24	0.25	0.3	0.74	0.61	0.22	0.52	0.5	0.26	0.25	0.3				
		葉素含有量	mg/L	0.013	0.032	0.035	0.025	0.03	0.023	0.025	0.026	0.038	0.03	0.032	0.067	0.026	0.026	0.04	0.032	0.026	0.03				
		りん含有量	mg/L	17.0	26.8																				